

## Influence of Polyphenol Extract from Evening Primrose (*Oenothera Paradoxa*) Seeds on Proliferation of Caco-2 Cells and on Expression, Synthesis and Activity of Matrix Metalloproteinases and Their Inhibitors

Karolina Szewczyk<sup>1\*</sup>, Urszula Lewandowska<sup>2</sup>, Katarzyna Owczarek<sup>2</sup>, Dorota Sosnowska<sup>3</sup>, Sylwia Goralach<sup>3</sup>, Maria Koziółkiewicz<sup>3</sup>, Zbigniew Hrabec<sup>1</sup>, Elżbieta Hrabec<sup>1</sup>

<sup>1</sup>Department of Medical Enzymology, Medical University of Lodz, Mazowiecka 6/8, 92-215 Lodz, Poland

<sup>2</sup>Department of Biochemistry, Medical University of Lodz, Mazowiecka 6/8, 92-215 Lodz, Poland

<sup>3</sup>Institute of Technical Biochemistry, Technical University of Lodz, Stefanowskiego 4/10, 90-924 Lodz, Poland

Key words: evening primrose, matrix metalloproteinases, cancer

Evening primrose (*Oenothera paradoxa* Hudziok) seeds are a rich source of not only a valuable oil containing an essential fatty acid -  $\gamma$ -linolenic acid (GLA) - but also polyphenols which can be obtained from the biomass remaining after oil pressing. The aim of our studies was to evaluate the influence of a polyphenol extract from defatted seeds of evening primrose on human colorectal adenocarcinoma Caco-2 cell proliferation and matrix metalloproteinases (MMPs) synthesis and activity. To assess the effect of evening primrose extract on Caco-2 cell proliferation, crystal violet staining and sulforhodamine B (SRB) assays were used whereas mRNA expression and activity of MMPs were evaluated by RT-PCR and gelatin zymography.

The results revealed that the examined polyphenol extract had little influence on Caco-2 proliferation, but effectively in a time- and dose-dependent manner inhibited *MMP-1*, *MMP-7*, *MMP-9* and *MMP-14* mRNA synthesis induced by  $\text{TNF-}\alpha$  and TPA. Additionally, zymographic analysis revealed that after 24 h, the polyphenol extract at a concentration of 50  $\mu\text{mol/L}$  GAE caused a 10-fold reduction in *MMP-9* synthesis. Moreover, this extract might be a potent inhibitor of MMP activity. The results showed that polyphenol extract from evening primrose inhibited PBMC-derived *MMP-2* and *MMP-9* enzymatic activity in dose-dependent manner. The obtained results indicate that the polyphenol extract from evening primrose seeds could be an inhibitor of proteases involved in tumor progression and metastasis.

### INTRODUCTION

Currently, cancer is the most frequent disease occurring in the modern world. Despite the progress in cancer research and treatment, as well as a huge amount of money given to find new, more effective drugs, it still remains a major worldwide health problem. Global statistics show that colorectal cancer ranked third place in males as the most commonly diagnosed cancer whereas in females ranked second [Jemal *et al.*, 2011]. Moreover, in Eastern Europe the incidence rate of colorectal cancer is still increasing [Center *et al.*, 2009].

Matrix metalloproteinases (MMPs) are endopeptidases containing  $\text{Zn}^{2+}$  in an active center and requiring  $\text{Ca}^{2+}$  for catalysis. On the basis of substrate specificity and domain organization MMPs are classified into: collagenases (*MMP-1*), gelatinases (also known as type IV collagenases: *MMP-2*, *MMP-9*), stromelysins (*MMP-3*), matrilysin (*MMP-7*) and membrane-type MMPs (*MMP-14*). Collectively they are able to degrade all protein components of the extracellular matrix (ECM) and some non-ECM proteins [Ra & Parks, 2007]. Under physiological conditions, the activities

of MMPs are precisely regulated; first of all, at the transcription level, by activation of the zymogens and by their natural inhibitors (tissue inhibitors of metalloproteinases; TIMPs). Except for neutrophils, cells do not accumulate MMPs. After synthesis, MMPs are secreted into the ECM as a latent form that can be activated by either proteolytic cleavage or by conformational changes induced, among others, by reactive oxygen species (ROS) [Hadler-Olsen *et al.*, 2013].

MMPs play a key role in physiological processes such as tissue repair and morphogenesis, as well as in many pathological processes including tumorigenesis, angiogenesis, metastasis [Kupai *et al.*, 2010]. These enzymes are produced by fibroblasts, monocytes, lymphocytes, neutrophils, endothelial cells as well as cancer cells. The cancer cell lines are the simplified models of tumor. *In vivo* cancer cells might be in contact with interleukins, growth factors, and cytokines, which induce MMP synthesis [Noël *et al.*, 2008]. Transcription factors  $\text{NF-}\kappa\text{B}$  and activator protein-1 (AP-1) are involved in the MAPK-mediated MMPs expression in response to pro-inflammatory cytokine  $\text{TNF-}\alpha$ . AP-1 can also be induced by phorbol-ester-12-*O*-tetradecanoylphorbol-13-acetate (TPA) [Labrie & St-Pierre, 2013]. Strategies for the prevention of proteolytic matrix degradation have mainly focused on MMPs. Synthetic MMP inhibitors such as bati-

\* Corresponding Author: Tel. +48 42 272 57 13; Fax: +48 42 272 56 79  
E-mail: [szewczyk\\_karolina@wp.pl](mailto:szewczyk_karolina@wp.pl) (K. Szewczyk)

mastat and marimastat, were developed and tested in clinical trials, first and foremost to treat cancer. However, the results from these trials in cancer patients were disappointing [Gialeli *et al.*, 2011].

As far as we know, highly effective anticancer drugs that could effectively inhibit MMP activities with minor side effects have not been found to date. Natural plant polyphenols are quite an interesting group of compounds that might have such properties.

In the past decade an increased interest in polyphenolic compounds occurring in our diet has been observed. Polyphenols are plant secondary metabolites widespread in the plant kingdom. They have many beneficial properties proved in *in vitro* and *in vivo* studies, namely: antibacterial [Daglia *et al.*, 2007], antifungal [Báidez *et al.*, 2006], neuroprotective [Kumar *et al.*, 2007] and also anticancer, in particular: antioxidant [Robaszekiewicz *et al.*, 2007], antimetastatic [Ogasawara *et al.*, 2007], proapoptotic [Mertens-Talcott & Percival, 2005], antiproliferative [Letchoumy *et al.*, 2007], antiangiogenic [Lamy *et al.*, 2006], anti-inflammatory [Clavin *et al.*, 2007]. More precisely, they can influence anti-apoptotic proteins (*e.g.* Bcl-2, Bcl-X<sub>L</sub>), proapoptotic proteins (*e.g.* caspases), cell cycle proteins (cyclins, cyclin-dependent kinases), protein kinases (*e.g.* IKK, EGFR, MAPK), cell adhesion molecules, growth factor signaling pathways and transcription factors (*e.g.* AP-1, STAT3) [extensively reviewed by González-Vallinas *et al.*, 2013]. The best-known polyphenols having well-documented anticancer activities are: epigallocatechin gallate (EGCG), resveratrol, isoflavons and derivatives of catechin. A number of reports suggest that these types of compounds not only may inhibit inflammation, which can lead to hyperproliferation and carcinogenesis, but also can inhibit angiogenesis and metastasis [González-Vallinas *et al.*, 2013].

Many herbal plants are under investigation for their potential as chemopreventive agents. In our studies we decided to examine a less known plant having health benefits. Evening primrose (*Oenothera paradoxa* Hudziok) is a weed originated from North America. In Europe it is cultivated because its seeds contain a high amount of polyunsaturated fatty acids, particularly  $\gamma$ -linolenic acid (GLA) [Peiretti *et al.*, 2004], which might be used as an alternative treatment of hypercholesterolemia or rheumatoid arthritis [Jaszewska *et al.*, 2010]. After extraction of oil the remaining defatted seeds contain polyphenolic compounds that can have beneficial health effect. So far, the extracts obtained from defatted seeds from *Oenothera paradoxa* Hudziok caused a decrease of plasma cholesterol and low density lipoprotein cholesterol in rats [Bałasińska, 1998], a dose-dependent inhibition of lipid peroxidation in mouse lymphocytic leukemic L1210 cells [Bałasińska & Troszyńska, 1998] a dose-dependent inhibition of metalloproteinases activity [Kiss *et al.*, 2008], induction of apoptosis in human skin melanoma HTB-140 cells [Jaszewska *et al.*, 2009] and reduction of cell viability and cell invasiveness in human breast cancer MDA-MB-231 cells [Lewandowska *et al.*, 2013].

We examined the influence of a polyphenol extract from defatted seeds of evening primrose (*Oenothera paradoxa*) on proliferation of human colorectal adenocarcinoma cell line (Caco-2) as well as their influence on the expression of several MMPs (especially *MMP-9*), their inhibitors (*TIMP-1*

and *TIMP-2*), and enzymatic activity of type IV collagenases (*MMP-2* and *MMP-9*) synthesized by human peripheral blood mononuclear (PBMC) cells.

## MATERIALS AND METHODS

### Plant materials

Evening primrose (*Oenothera paradoxa* Hudziok) defatted seeds were obtained from Agropharm S.A./Adamed Group pharmaceutical company (Tuszyn, Poland).

### Chemicals

All the reagents, cell culture medium and its supplements were purchased from Sigma-Aldrich Co. LLC (St. Louis, MO), except for penicillin, streptomycin – Polfa Tarchomin (Tarchomin, Poland) and amphotericin B – Biochrom AG (Berlin, Germany). TRIzol Reagent, oligo (dT)<sub>20</sub>, random hexamer and SuperScript™ II Reverse Transcriptase were obtained from Life Technologies Corp. (Eugene, OR). Reagents for PCR reaction were bought from Finnzymes OY (Espoo, Finland). Cryptochlorogenic acid, procyanidin B1, procyanidin B2 and procyanidin C1 were purchased from PhytoLab GmbH & Co.KG (Vestenbergsgreuth, Germany). HPLC grade acetonitrile was purchased from J.T. Baker (Griesheim, Germany).

### Polyphenol extract preparation

Evening primrose dry polyphenol extract was obtained according to the method described by Goralch *et al.* [2011]. Briefly, the waste defatted seeds obtained from Agropharm S.A./Adamed Group pharmaceutical company were milled and defatted with hexane.

Then the seeds underwent triple extraction with 70% aqueous solution of ethanol. The obtained extracts were centrifuged (4000 rpm, 15 min) and then concentrated under vacuum at the temperature below 40°C. The obtained aqueous solution of the polyphenols was lyophilized. The dry extract was stored at –20 °C prior to further analyses.

### Characterization of the polyphenol extract

The extract was characterized in terms of total polyphenol content by the Folin-Ciocalteu procedure, expressed as (+) catechin equivalents [Peri & Pompei, 1971], total flavanol content by the vanillin procedure, expressed as (+)catechin equivalents [Swain & Hillis, 1959], and total proanthocyanidin content after acid hydrolysis in butanol environment, expressed as cyanidin [Rösch *et al.*, 2003]. Ellagitannins and gallotannins (after acid hydrolysis in methanol environment, 20 h, 85°C) were determined by an analytical reversed-phase HPLC system with the use of a Eurospher-100 C18 column (250 mm x 4.6 mm, 5 mm) (Knauer, Berlin, Germany). A binary mobile phase and a gradient program were the same as described below. Ellagitannin content determined at 254 nm is expressed as ellagic acid and gallotannin content at 280 nm is expressed as methyl gallate [Hartzfeld *et al.*, 2002].

### Phenolics determination by HPLC-DAD

Phenolic profile was determined using an analytical reversed-phase HPLC system (Waters, Milford, MA) with

a 2707 autosampler and a 1525 binary HPLC pump coupled to a 996 photodiode array detector (2998), controlled by Waters Breeze 2 software (Waters). A SYMMETRY C18 column (250 mm x 4.6 mm, 5 mm) (Waters) was used. According to Dyrby *et al.* [2001], the binary mobile phase consisted of water and formic acid in the ratio of 90:10 (v/v), respectively (solvent A); water, acetonitrile and formic acid in the ratio of 49:50:10 (v/v/v), respectively (solvent B). The phenolic separation was performed using the following gradient program with a flow rate of 1 mL/min: 0 min, 88% A + 12% B; 26 min, 70% A + 30% B; 40–43 min, 0% A + 100% B; 48–50 min, 88% A + 12% B. Detection was performed by scanning from 200 to 550 nm. Peak identification was carried out by comparison of retention times and diode array spectral characteristics with the standards.

### Cell culture

A human colorectal adenocarcinoma cell line (Caco-2) was purchased from the Institute of Immunology and Experimental Therapy, Polish Academy of Sciences (Wroclaw, Poland). Caco-2 cells were cultured in MegaCell™ MEM culture medium supplemented with 3% heat-inactivated fetal bovine serum (FBS), 4 mmol/L L-glutamine, 100 U/mL penicillin, 100 µg/mL streptomycin and 1.25 µg/mL amphotericin B at 37°C in a humidified atmosphere containing 5% CO<sub>2</sub>. All experiments were carried out between passages 3 and 19. The cells were seeded in such quantity that the confluence at the end of the experiment did not exceed 80% in control wells. Cell viability was assessed on the basis of trypan blue exclusion.

The polyphenol extract was tested at concentrations ranging from 25 to 100 µmol/L gallic acid equivalents (GAE), which corresponds to 4.3–17.0 µg GAE/mL or 7.3–29.0 µg extract/mL. The solution of the preparation was made first in 70% ethanol and then diluted to the final concentration with a serum-free medium. The final concentration of ethanol was lower than 0.01% (v/v). The polyphenol content in the extract is expressed as gallic acid equivalents since it represents the mean value between (+)catechin equivalents and ellagic acid equivalents.

### Crystal violet staining

To assess the effect of evening primrose extract on the proliferation of Caco-2 crystal violet staining and sulforhodamine B (SRB) assay were used. Crystal violet staining (CVS) was performed according to Henriksson *et al.* [2006] with slight modifications. Briefly, the cells were harvested, suspended in the growth medium mentioned above and seeded on 96-well plates. After 24 h, the cells were exposed to the polyphenol extract from evening primrose or EGCG or gallic acid for the following 24, 48 and 72 h. After culture, the medium was removed and the cells were fixed *in situ* with 4% formaldehyde in PBS (phosphate buffered saline) for 30 min at room temperature. Then, the cells were washed twice with PBS and stained with 0.5% crystal violet in 25% methanol aqueous solution for 5 min at room temperature. Unbound dye was washed out with deionized water and the cells were allowed to air-dry. The dye was dissolved in 33% acetic acid aqueous solution on a shaker for 30 min at room temperature. OD was measured by a 96-well plate reader (iMark™, BioRad, Herkules, CA, USA) at the wavelength of 595 nm.

### Sulforhodamine B assay

Sulforhodamine is an anionic dye, which migrates into the cell and binds to basic amino acid residues of proteins under mild acidic conditions [Houghton *et al.*, 2007]. Briefly, the cells were harvested, suspended in the growth medium mentioned above and seeded on 96-well plates. After 24 h, the cells were exposed to the polyphenol extract from evening primrose or EGCG or gallic acid for the following 24, 48 and 72 h. Next, the culture medium was removed and the cells were fixed *in situ* with ice-cold 10% aqueous solution of trichloroacetic acid (TCA) for 1 h at 4°C. After that, TCA was removed and each well was washed with deionized water and allowed to air-dry. The cells were stained for 30 min at room temperature with 0.4% SRB solution in 1% aqueous solution of acetic acid. The dye excess was removed and each well was washed with 1% acetic acid and allowed to air-dry. The protein-bound dye was solubilized by 10 mmol/L unbuffered Tris on a shaker for 10 min at room temperature. The optical density (OD) was measured by a 96-well plate reader (iMark™, BioRad, Herkules, CA, USA) at the wavelength of 490 nm.

### Gelatin zymography and quantitative analysis of MMP-9 secretion

Caco-2 cells were harvested, suspended in the growth medium mentioned above and seeded on 96-well plates. After 24 h, the cells were washed twice with PBS and then suspended in the medium without FBS. Subsequently, the cells were treated with 10 ng/mL tumor necrosis factor α (TNF-α), 10 ng/mL 12-*O*-tetradecanoylphorbol-13-acetate (TPA), the polyphenol extract from evening primrose or gallic acid or EGCG and incubated for 24 and 48 h. MMP-9 is an inducible enzyme, therefore TNF-α and TPA were added in order to stimulate Caco-2 for its synthesis. Afterwards, conditioned media were harvested and stored at –20°C for further analysis. MMP-9 secretion was determined by means of zymographic analysis. Briefly, the same volumes of the conditioned media (usually 20 µL) were mixed with a sample buffer containing 4% SDS (sodium dodecyl sulfate), 30% glycerol, 0.25 mol/L Tris-HCl (pH 6.8) and 0.01% bromophenol blue. Electrophoresis was carried out in 10% polyacrylamide gel containing 1 mg/mL gelatin. After the electrophoresis, SDS was removed from the gels by incubation in a buffer containing 2% Triton X-100 and 50 mmol/L Tris-HCl (pH 7.4) and the enzyme reaction was allowed to proceed at 37°C for 21 h in a buffer containing 1% Triton X-100, 50 mmol/L Tris-HCl (pH 7.4) and 5 mmol/L CaCl<sub>2</sub>. Next, the gels were stained for 1 h in a solution containing 0.1% amido black, 7% acetic acid and 20% ethanol. After the staining procedure, the gels were captured using an Olympus camera (Olympus Corp., Tokyo, Japan). MMP-9 was visualized as a transparent band against the dark blue background of the amido black-stained gels. Densitometry analysis was carried out using Quantity One® 4.4 software (BioRad, Herkules, CA).

### Reverse Transcriptase – Polymerase Chain Reaction (RT-PCR)

The cells were harvested, suspended in the growth medium mentioned above and seeded on culture dishes. After 24 h, the cells were washed twice with PBS and then suspended

TABLE 1. Sequences of the examined primers.

Gene	Primer sequence 5'→3'	Annealing temperature	Product size
<i>MMP-1</i>	L: AGCTCAGGATGACATTGATGG R: AGCTCAACTTCCGGGTAGAAG	50°C	198 bp
<i>MMP-7</i>	L: TCAGGAAAGTTGTATGGGGAAC R: GCTAAATGGAGTGGAGGAACAG	50°C	471 bp
<i>MMP-9</i>	L: ACTTTGACAGCGACAAGAAGTG R: CCCTCAGTGAAGCGGTACATAG	49°C	151 bp
<i>MMP-14</i>	L: CATCCAGAAGAGAGCAGCATC R: GAGAGGAAGGATGGCAAATTC	54°C	150 bp
<i>TIMP-1</i>	L: AGCCAACAGTGTAGGTCTTGG R: TACTTCCACAGGTCCCACAAC	49°C	157 bp
<i>TIMP-2</i>	L: TTCTCTGTGACCCAGTCCATC R: ACCCTCTGTGACTTCATCGTG	49°C	170 bp
<i>GAPDH</i>	L: TGATGACATCAAGAAGGTGGTGAAG R: TCCTTGAGGCCATGTGGGCCAT	54°C	240 bp

ed in the medium without FBS. Subsequently, the cells were exposed to 10 ng/mL TNF- $\alpha$ , 10 ng/mL TPA and the evening primrose polyphenol extract for 16 h. After incubation, the cells were lysed by the TRIzol reagent (1 mL per 10<sup>6</sup> cells). Further, RNA isolation was performed according to the manufacturer's protocol. The obtained RNA pellet was allowed to air-dry and then was redissolved in RNase-free water. Total RNA content was quantified using a Beckman DU650 spectrophotometer (Beckman Coulter, Inc., Brea, CA) at 260 nm. The samples were stored at -80°C for further analysis.

cDNA was synthesized using total RNA template, oligo (dT)<sub>20</sub>, random hexamer and SuperScript™ II Reverse Transcriptase (RT). The reaction was carried out according to the manufacturer's protocol enclosed with SuperScript™ II RT. Then the samples were stored at -80°C for further analysis.

The cDNA amplification was carried out in 50  $\mu$ L volume. The reaction mixture consisted of: 5  $\mu$ L cDNA template, DyNAzyme EXT buffer (50 mmol/L Tris-HCl, pH 9.0, 1.5 mmol/L MgCl<sub>2</sub>, 15 mmol/L (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0.1 % Triton X-100), 10  $\mu$ mol/L dNTPs, 1 U DyNAzyme™ EXT DNA Polymerase and 10 pmol forward and reverse primers (presented in Table 1). The reaction was submitted to 35 cycles of denaturation (94°C for 30 sec), annealing (for 30 sec at temperatures given in Table 1) and extension (72°C for 60 sec). The amplified products were detected in 2% agarose gel containing 0.5  $\mu$ g/mL ethidium bromide, documented by VersaDoc™ Imaging System (BioRad, Hercules, CA) and analyzed using Quantity One® 4.4 software (BioRad). The amount of RT-PCR product for the gene of interest was normalized to the amount of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) in the same sample.

#### Isolation and culture of human peripheral blood mononuclear cells (PBMC), and zymographic detection of PBMC-derived type IV collagenases after incubation with the evening primrose polyphenol extract

Peripheral blood collected from a healthy volunteer donor was diluted with PBS in the ratio of 1:1 and layered onto Histopaque-1077. After centrifugation at 400 $\times$ g for 30 min at

room temperature, the layer of mononuclear cells was collected, and the cells were washed twice with PBS and suspended in RPMI 1640 culture medium containing 25 mmol/L HEPES, 2 mmol/L L-glutamine, 100 U/mL penicillin, 100  $\mu$ g/mL streptomycin, 1.25  $\mu$ g/mL fungizone and 10% FBS. The cells were cultured at 37°C in a humidified atmosphere of 5% CO<sub>2</sub> at the initial concentration of 1 $\times$ 10<sup>6</sup> per 1 mL. PBMC-conditioned medium was used as a source of MMP-2 and MMP-9. Zymographic detection of PBMC-derived type IV collagenases after incubation with the evening primrose seed extract was carried out according to a procedure described previously [Stręk *et al.*, 2007]. For the zymographic assay, the extract was dissolved in 70% ethanol aqueous solution and further diluted with the incubation buffer so that the final concentration of ethanol was <0.01% (v/v).

#### Statistical analysis

Statistical comparisons were made using nonparametric Kruskal-Wallis test and one-way ANOVA followed by Bonferroni test (Analyze-it software, v. 2.21, Leeds, UK). Data were presented as mean  $\pm$  SEM. A *p* value of less than 0.05 was considered to be significant.

## RESULTS

#### Characterization of the polyphenol extract from evening primrose seeds

The examined extract was isolated according to the procedure presented above. The characterization of the preparation is presented in Table 2. HPLC analysis revealed that the evening primrose polyphenol extract contained among others: gallic acid, procyanidin B1, procyanidin B2, (+)-catechin, procyanidin C1, (-)-epicatechin and ellagic acid.

#### Influence of the polyphenol preparation on Caco-2 proliferation (CVS, SRB assay)

The influence of the polyphenol extract from evening primrose on the proliferation of Caco-2 was evaluated color-

TABLE 2. Characterization of the polyphenol extract from defatted evening primrose seeds.

Compound	Content (mg/g of dry extract)
Total polyphenols <sup>a</sup>	668.71 ± 15.76
Total flavanols <sup>b</sup>	190.02 ± 5.32
Total proanthocyanidins <sup>c</sup>	142.86 ± 4.05
Total flavanols and HBA <sup>d</sup>	n.a.
Total flavanols and HBA <sup>e</sup>	228.05 ± 4.65
(+)catechin	11.77 ± 1.24
(-)-epicatechin	11.36 ± 0.87
Procyanidin B1	7.34 ± 0.36
Procyanidin B2	19.47 ± 0.95
Procyanidin C1	7.84 ± 1.24
Chlorogenic acid	n.d.
Cryptochlorogenic acid	n.d.
Other HCA acids <sup>f</sup>	0.16 ± 0.01
Quercetin 3-rhamnoside	n.d.
Other flavonols <sup>g</sup>	0.78 ± 0.05
Free ellagic acid	2.65 ± 0.23
Free gallic acid	5.17 ± 0.11
Ellagitannins <sup>h</sup>	4.12 ± 0.19
Gallotannins <sup>i</sup>	89.08 ± 8.13

Mean ± SD,  $n \geq 3$ . <sup>a</sup>determined by Folin-Ciocalteu reagent as (+) catechin equivalents. <sup>b</sup>determined by vanillin reagent as (+)catechin equivalents. <sup>c</sup>determined by acid hydrolysis as cyanidin equivalents. <sup>d</sup>determined by HPLC at 280 nm as (+)catechin. <sup>e</sup>determined by HPLC at 280 nm as gallic acid. <sup>f</sup>determined by HPLC at 320 nm as chlorogenic acid. <sup>g</sup>determined by HPLC at 360 nm as quercetin. <sup>h</sup>determined by HPLC at 254 nm by acid hydrolysis as ellagic acid. <sup>i</sup>determined by HPLC at 280 nm by acid hydrolysis as methyl gallate. HBA - hydroxybenzoic acids; HCA - hydroxycinnamic acids; n.d. - not detected; n.a. - not analyzed.

metrically after staining the cells with crystal violet and sulforhodamine B. The cells were seeded on plates and exposed to polyphenols. After the exposure time, the number of cells was evaluated according to the protocols described above. The analysis revealed that 25  $\mu\text{mol/L}$  GAE of the evening primrose polyphenol extract seemed to stimulate the proliferation of Caco-2 (especially after 24 and 48 h incubation) while this effect was not observed at higher concentrations (Figure 1A). However, after 72 h one can observe a decline in the cell number at 100  $\mu\text{mol/L}$  GAE.  $\text{IC}_{50}$  was not reached in the experiment. EGCG, the main green tea polyphenol with well-documented antiproliferative effect, at the concentration of 25  $\mu\text{mol/L}$  stimulates the cell proliferation but higher concentrations (50, 75 and 100  $\mu\text{mol/L}$ ) suppress their division (Figure 1B). What is worth mentioning is the fact that gallic acid, one of the major constituents of polyphenol extract, stimulates Caco-2 proliferation at each concentration tested (Figure 1C). The lack of the antiproliferative effect of the polyphenol extract from evening primrose on Caco-2 might be caused by the high content of gallic acid in this preparation. The same tendency was observed when SRB was used for proliferation assay (data not shown).




FIGURE 1. The influence of the examined polyphenols: A) the evening primrose extract expressed as gallic acid equivalents (GAE), B) EGCG and C) gallic acid on Caco-2 cell number determined by crystal violet staining (CVS). Data represent the mean ± SEM of three independent experiments. Statistical significance of differences between means: \*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$  vs. control.

### Influence of the evening primrose polyphenol extract on expression of selected MMPs and their inhibitors

In the present work we assessed the influence of the polyphenol preparation from evening primrose on the expression of *MMP-1*, *MMP-7*, *MMP-9*, *MMP-14*, *TIMP-1* and *TIMP-2*. In our experiment a low level of constitutive synthesis of *MMP-9* was detectable in untreated Caco-2 cells but the expression of this gene can be stimulated with  $\text{TNF-}\alpha$  and TPA (Figure 2). The expression of *MMP-1* was not detected in cells not treated with  $\text{TNF-}\alpha$  and TPA, but the addition of these two compounds induced the expression of this protease (Figure 3). In our experimental conditions,  $\text{TNF-}\alpha$  and TPA stimulation of Caco-2 did not influence the expression levels of *MMP-7*, *MMP-14*, *TIMP-1* and *TIMP-2*.

Our studies revealed that the polyphenol extract from evening primrose inhibited the expression of *MMP-1*, *MMP-7*, *MMP-9* and *MMP-14* in a dose-dependent manner. Among the MMPs investigated, the polyphenol extract seems to be the most potent inhibitor for *MMP-7*, *MMP-9* and *MMP-14*.




FIGURE 2. The inhibitory effect of the polyphenol extract from evening primrose on the expression of *MMP-9* determined by RT-PCR. The concentrations of the extract are expressed as gallic acid equivalents (GAE). Data represent the mean  $\pm$  SEM of three independent experiments. Statistical significance of differences between means: \*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$  vs. control.

The preparation did not significantly influence the expression of *TIMP-1* and *TIMP-2* (data not shown).

### Gelatin zymography and quantitative analysis of MMP-9 secretion

In our study, MMP-9 expression was induced in Caco-2 cells by TNF- $\alpha$  and TPA as non-stimulated cells do not secrete this enzyme to the medium (Figure 4, lane 2). Therefore, to determine the influence of the polyphenol extract of defatted evening primrose seeds on MMP-9 synthesis and secretion, the cells were stimulated with inducers. Gelatin zymography can also be used to detect the presence of MMP-2 (the enzyme produced mainly by fibroblasts in tumor environment which is responsible for tissue remodelling and tumor progression). However, MMP-2 was not detected in Caco-2.

Caco-2 cells were incubated with the polyphenol preparation at different concentrations. After experiment, the conditioned media were collected and subjected to zymographic analysis. The results revealed that the polyphenol extract inhibited the synthesis and the secretion of MMP-9 in a time- and dose-dependent manner (Figure 4A). The polyphenol extract was a more potent inhibitor of MMP-9 secretion than EGCG as it is presented in Figure 4. One of the major constituents of the examined extract – gallic acid – did not influence the MMP-9 secretion (Figure 4C).

### Inhibition of MMP-2 and MMP-9 activities by the evening primrose polyphenol extract

PBMC secrete both type IV collagenases in approximately equal amounts, as shown in Figure 5 and by Stręk *et al.* [2007]. We therefore decided to use PBMC-conditioned medium as a source of MMP-2 and MMP-9 for our study on the influence of the *O. paradoxa* defatted seed extract on the activities of those enzymes. The extract inhibited the activities of both type IV collagenases in a concentration-depen-




FIGURE 3. The influence of the examined polyphenol extract on the expression of: A) *MMP-1*, B) *MMP-7*, C) *MMP-14* in Caco-2 cell line determined by RT-PCR. The concentrations of the extract are expressed as gallic acid equivalents (GAE). Data represent the mean  $\pm$  SEM of three independent experiments. Statistical significance of differences between means: \*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$  vs. control.

dent manner. As estimated from Figure 5,  $IC_{50}$  was between 7 and 22  $\mu\text{mol/L}$  GAE for both enzymes, and their inhibition levels were similar to each other at all concentrations tested.

## DISCUSSION

The oil from the seeds of *Oenothera paradoxa* is the best known dietary supplement obtained from this plant; however, there are few studies dealing with polyphenol extracts from defatted seeds of evening primrose [Jaszewska *et al.*, 2009; Górlach *et al.*, 2011]. In our studies, we decided to explore some biological properties of this polyphenol extract.

The polyphenol extract was subjected to HPLC analysis. In this preparation we were able to identify: gallic acid, procyanidin B1, procyanidin B2, (+)catechin, procyanidin C1, (-)epicatechin and ellagic acid. Additionally, Górlach




FIGURE 4. Zymographic analysis of TNF- $\alpha$ - and TPA-induced MMP-9 secretion by Caco-2 after 24 and 48 h incubation with: A) the evening primrose polyphenol extract expressed as gallic acid equivalents (GAE), B) EGCG and C) gallic acid. The right panel presents zymograms of conditioned media after 48 h as an example of polyphenols inhibitory action. Data represent the mean  $\pm$  SEM of three independent experiments. Statistical significance of differences between means: \*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$  vs. control.




FIGURE 5. The influence of the evening primrose polyphenol extract on MMP-2 and MMP-9 activities, assessed by gelatin zymography. The concentrations of the extract are expressed as gallic acid equivalents (GAE). Data are presented as the mean  $\pm$  SEM, of three independent experiments. Statistical significance of differences between means: \*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$  vs. control.

*et al.* [2011] using MALDI-TOF MS confirmed the presence of (+)catechin, (-)epicatechin, B-type dimeric procyanidin digallate and/or B-type dimer of (epi)catechin gallate, A-type procyanidin dimer, B-type procyanidin trimer, procyanidin tetramer and hydrolysable tannins (mono- and trigalloylglucose). Kiss *et al.* [2008] analyzed the composition of a polyphenol extract prepared from evening primrose defatted seeds originated from the same source as in our study (Agropharm S.A./Adamed Group, Tuszyn, Poland) and were also able to identify: (-)epicatechin gallate, procyanidin B3 (dimer), oenothien B (macrocylic ellagitannin dimer) and penta-*O*-galloyl- $\beta$ -D-glucose (PGG), which indicates that the polyphenol extract from evening primrose is a reservoir of versatile biological compounds.

The aim of our study was to evaluate the effect of the polyphenol extract from defatted seeds of evening primrose on human colorectal adenocarcinoma cell line (Caco-2).

We investigated the effect of the polyphenol extract on cell growth by means of crystal violet staining and sulforhodamine B assay. We intentionally did not use 3-[4,5-dimethyl-

thiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT) assay in our experiments, as Bernhard *et al.* [2003] revealed that resveratrol increased MTT-reducing activity without an observed increase in cell number showing that this test is ineffective in the evaluation of cell proliferation after treatment with polyphenolic compounds. Therefore, to determine cell growth we have chosen such reagents that do not interfere with phenolic compounds. In our studies, the proliferation assays (CVS and SRB) revealed that the preparation from evening primrose exhibited a stimulating effect on Caco-2 proliferation at low concentrations whereas at higher concentrations had rather no influence on cell division. Gallic acid, one of the components of this preparation, exhibited a similar effect on Caco-2 cells. Other studies showed that the influence of the polyphenol extract on proliferation of cancer cells might be a cell-specific phenomenon. For instance, Li *et al.* [2010] testing a polyphenol extract from Devil's Club (*Oplopanax horridus*) on different colorectal cancer cell lines observed that this preparation did not influence HT-29 proliferation at all whereas significantly reduced the growth of HCT-116 cell line. Moreover, in our studies, we observed a strong inhibition of MDA-MB-231 human breast cancer cell division by the polyphenol extract from evening primrose [Lewandowska *et al.*, 2013].

It is well documented that MMPs, especially type IV collagenases, play a key role in the development and progression of cancer [Kessenbrock *et al.*, 2010]. Therefore, we decided to investigate whether this preparation might influence the expression of matrix metalloproteinase 9. Augmented MMP-9 expression frequently correlates with cancer progression and metastasis [Kessenbrock *et al.*, 2010], as the enzyme is able to degrade components of ECM, especially collagen type IV, the main component of the basement membranes. High expression of *MMP-9* was observed in colorectal cancer cells as well as in stromal cells [Lubbe *et al.*, 2006]. Moreover, in animal studies where human pancreatic cancer cells were implanted into nude mice, a rapid growth of tumor cells was observed in the mice expressing *MMP-9* gene, whereas the animals lacking this gene produced significantly smaller tumors [Nakamura *et al.*, 2007]. In our studies, we induced the expression of *MMP-9* in Caco-2, as in this cancer cell line the *MMP-9* expression was on the very low level. *MMP-9* was both TNF- $\alpha$ - and TPA-induced, as applying only one of the stimuli was insufficient to obtain the satisfying results (data not shown). Probably, the up-regulation of *MMP-9* expression in these cells requires binding of two activated transcription factors, namely AP-1 and NF- $\kappa$ B, to the transcription factor binding motifs in *MMP-9* promoter region at the same time [Labrie & St-Pierre, 2013]. There is probably a synergistic effect of both transcription factors.

To evaluate the influence of the polyphenol extract on *MMP-9* expression, the cells were seeded on culture plates and the extract at different concentrations was added simultaneously with TNF- $\alpha$  and TPA. After 16 h cells were lysed, and total RNA was extracted and used for RT-PCR assays. In parallel with this experiment the cells were cultured for 48 h and the conditioned media were subjected to zymographic analysis.

The results have shown that the polyphenol extract from evening primrose has a high potency to inhibit *MMP-9* expres-

sion, as assessed by RT-PCR and gelatin zymography. Zymographic analysis revealed that the polyphenols from evening primrose were able to inhibit *MMP-9* secretion even more effectively than EGCG. These data suggest that the preparation might significantly attenuate tumor progression by blocking *MMP-9* synthesis. The main polyphenol from green tea, EGCG, likewise the polyphenols tested, was able to decrease *MMP-9* expression in 95-D lung carcinoma cells at the concentration of 40  $\mu$ mol/L [Yang *et al.*, 2005], whereas in periodontal diseases it was effective even at the concentration of only 20  $\mu$ mol/L [Yun *et al.*, 2004].

Other matrix metalloproteinases, such as *MMP-1* and *MMP-7*, similarly to *MMP-9* exhibit proteolytic activity towards ECM, and these two enzymes also correlate with an aggressive phenotype of colorectal cancer [Hadler-Olsen *et al.*, 2013]. Our study is the first report on the inhibition of *MMP-1* expression by the evening primrose polyphenol extract in a colon cancer cell line. Similar effects were observed by Toegel *et al.* [2012], who examined the influence of a *Caesalpinia sappan* extract on human bone chondrosarcoma SW1353 cells. What is worth mentioning is that contrary to the results of Kim *et al.* [2007], who reported that EGCG stimulates the expression of *MMP-7* in colorectal cancer cell line HT-29, our polyphenol extract proved to be an effective repressor of *MMP-7* gene. The activity of *MMP-7* is thought to promote tumor cell survival by shedding Fas receptor from the cell surface which is responsible for apoptosis of the cell [Li *et al.*, 2006].

*MMP-14* (membrane-type(MT)1-MMP) is one of the six membrane-bound MMPs which also plays a significant role in tumor progression by activating other members of the MMP family participating in cancer development, especially *MMP-2* [Poincloux *et al.*, 2009]. In our studies, we observed a strong inhibition of *MMP-14* expression by the polyphenol extract from evening primrose. These results indicate that the polyphenols tested effectively lowered *MMP-14* mRNA level, in contrast with the results obtained by Huang *et al.* [2011a], for an extract from *Phyllanthus urinaria* which did not change the level of *MMP-14* mRNA in human umbilical vein endothelial cells (HUVECs).

MMPs activity is regulated, among others, by the natural inhibitors – TIMPs (Tissue Inhibitors of Matrix Metalloproteinases). To date, four proteins belonging to this family (*TIMP-1-4*) have been described [Brew & Nagase, 2010]. The expression of tissue inhibitors of matrix metalloproteinases, namely *TIMP-1* and *TIMP-2*, was unaffected by the examined extract (data not shown).

The evening primrose polyphenol extract tested in the present work not only reduced *MMP-1*, -7, -9 and -14 expression in Caco-2 cells (as discussed above), but also inhibited the activities of PBMC-derived *MMP-2* and -9 in a concentration-dependent manner. There are literature data showing the inhibitory activity of evening primrose seed polyphenol-rich extracts obtained with the use of several solvents against two metalloproteinases, namely aminopeptidase N and neutral endopeptidase [Kiss *et al.*, 2008, 2012]. The highest inhibition was observed for aminopeptidase N ( $IC_{50} = 2.8$  and  $2.9 \mu$ g/mL for aqueous and 30% isopropanolic extracts, respectively) [Kiss *et al.*, 2008]. However, to



the best of our knowledge, the present work is the first report on type IV collagenase activity inhibition by a polyphenol-rich extract derived from evening primrose seeds. The inhibition mechanism of MMPs activity is still unknown. Ellagic acid, also the constituent of investigated polyphenol extract, was reported as an MMP-2 inhibitor with zinc-chelating properties exerted *in situ*, i.e. in the active center [Huang *et al.*, 2011b]. The authors suggested that a complex is formed in MMP-2 active center as a result of ellagic acid binding to the catalytic Zn-coordinated water molecule by its phenol group. According to Kołomecki [2000], blocking substrate cleavage by MMPs is one of the strategies of their inhibition. The inhibition of MMP activities observed on zymographic bands after their incubation with the evening primrose polyphenol extract may result from binding of high MW tannins to type IV collagen and, consequently, blocking the access of type IV collagenases to their substrate. The research conducted by Garbisa *et al.* [2001] investigating the inhibition of MMP-2 and -9 activities revealed that EGCG – a low molecular weight (MW) compound – exhibited a higher affinity towards type IV collagenases than to their substrate (gelatin). On the other hand, high MW tannins exhibited much stronger affinity to proteins having an open conformation (such as collagen and its denaturation product, gelatin) than to globular proteins [Naczek *et al.*, 2001; Deaville *et al.*, 2007].

## CONCLUSION

To sum up, the polyphenol extract from defatted seeds of evening primrose might modulate MMP-9 expression and secretion in a dose- and time-dependent manner. The presented results also indicate that it influences the expression of other matrix metalloproteinases (*MMP-1*, *MMP-7* and *MMP-14*) which are involved in cancer progression. Furthermore, the extract inhibited the activities of both type IV collagenases. Therefore, our preparation might significantly reduce the metastatic potential of colon cancer cells, however, further studies are needed.

## ACKNOWLEDGEMENTS

We would like to kindly thank Dorota Kustra, MSc for preparation of the polyphenol extract. This work was supported in part by grants from the Polish Ministry of Science and Higher Education (No.: N312 028 32/1893 and 4468/B/PO1/2011/40) and by a scholarship supporting PhD students' innovative researches, co-funded from the European Social Fund and the National Budget.

## REFERENCES

- Báidez A.G., Gómez P., Del Río J.A., Ortuño A., Antifungal capacity of major phenolic compounds of *Olea europaea* L. against *Phytophthora megasperma* Drechsler and *Cylindrocarpum destructans* (Zinssm.) Scholten. *Physiol. Mol. Plant Pathol.*, 2006, 69, 224–229.
- Bałańska B., Hypocholesterolemic effect of dietary evening primrose (*Oenothera paradoxa*) cake extract in rats. *Food Chem.*, 1998, 63, 453–459.
- Bałańska B., Troszyńska A., Total antioxidative activity of evening primrose (*Oenothera paradoxa*) cake extract measured *in vitro* by liposome model and murine L1210 cells. *J. Agric. Food Chem.*, 1998, 46, 3558–3563.
- Bernhard D., Schwaiger W., Crazzolara R., Tinhofer I., Kofler R., Csordas A., Enhanced MTT-reducing activity under growth inhibition by resveratrol in CEM-C7H2 lymphocytic leukemia cells. *Cancer Lett.*, 2003, 195, 193–199.
- Brew K., Nagase H., The tissue inhibitors of metalloproteinases (TIMPs): an ancient family with structural and functional diversity. *Biochim. Biophys. Acta*, 2010, 1803, SI, 55–71.
- Center M.M., Jemal A., Smith R.A., Ward E., Worldwide variations in colorectal cancer. *CA Cancer J. Clin.*, 2009, 59, 366–378.
- Clavin M., Gorzalczyński S., Macho A., Muñoz E., Ferraro G., Acevedo C., Martino V., Anti-inflammatory activity of flavonoids from *Eupatorium arnottianum*. *J. Ethnopharmacol.*, 2007, 112, 585–589.
- Daglia M., Papetti A., Grisoli P., Aceti C., Dacarro C., Gazzani G., Antibacterial activity of red and white wine against oral streptococci. *J. Agric. Food Chem.*, 2007, 55, 5038–5042.
- Deaville E.R., Green R.J., Mueller-Harvey I., Willoughby I., Frazier R.A., Hydrolyzable tannin structures influence relative globular and random coil protein binding strengths. *J. Agric. Food Chem.*, 2007, 55, 4554–4561.
- Dyrby M., Westergaard N., Stapelfeldt H., Light and heat sensitivity of red cabbage extract in soft drink model systems. *Food Chem.*, 2001, 72, 431–437.
- Garbisa S., Sartor L., Biggin S., Salvato B., Benelli R., Albin A., Tumor gelatinases and invasion inhibited by the green tea flavanol epigallocatechin-3-gallate. *Cancer*, 2001, 91, 822–832.
- Gialeli C., Theocharis A.D., Karamanos N.K., Roles of matrix metalloproteinases in cancer progression and their pharmacological targeting. *FEBS J.*, 2011, 278, 16–27.
- González-Vallinas M., González-Castejón M., Rodríguez-Casado A., Ramírez de Molina A., Dietary phytochemicals in cancer prevention and therapy: a complementary approach with promising perspectives. *Nutr. Rev.*, 2013, 71, 585–599.
- Gorlach S., Wagner W., Podsedek A., Sosnowska D., Dastych J., Koziolkiewicz M., Polyphenols from evening primrose (*Oenothera paradoxa*) defatted seeds induce apoptosis in human colon cancer Caco-2 cells. *J. Agric. Food Chem.*, 2011, 59, 6985–6997.
- Hadler-Olsen E., Winberg J.O., Uhlin-Hansen L., Matrix metalloproteinases in cancer: their value as diagnostic and prognostic markers and therapeutic targets. *Tumor Biol.*, 2013, 34, 2041–2051.
- Hartzfeld P.W., Forkner R., Hunter M.D., Hagerman A.E., Determination of hydrolyzable tannins (gallotannins and ellagitannins) after reaction with potassium iodate. *J. Agric. Food Chem.*, 2002, 50, 1785–1790.
- Henriksson E., Kjellén E., Wahlberg P., Wennerberg J., Kjellström J.H., Differences in estimates of cisplatin-induced cell kill in vitro between colorimetric and cell count/colony assays. *In Vitro Cell Dev. Biol. Anim.*, 2006, 42, 320–323.
- Houghton P., Fang R., Techatanawat I., Steventon G., Hylands P.J., Lee C.C., The sulphorhodamine (SRB) assay and other approaches to testing plant extracts and derived compounds for activities related to reputed anticancer activity. *Methods*, 2007, 42, 377–387.
- Huang S.T., Wang C.Y., Yang R.C., Wu H.T., Yang S.H., Cheng Y.C., Pang J.H., Ellagic acid, the active compound of *Phyllanthus*

- urinaria*, exerts *in vivo* anti-angiogenic effect and inhibits MMP-2 activity. *Evid-Based Compl. Altern. Med.*, 2011a, DOI: 10.1093/ecam/nep207.
20. Huang S.T., Yang R.C., Wu H.T., Wang C.N., Pang J.H., Zinc-chelation contributes to the anti-angiogenic effect of ellagic acid on inhibiting MMP-2 activity, cell migration and tube formation. *PLoS One*, 2011b, 6(5), e18986.
  21. Ii M., Yamamoto H., Adachi Y., Maruyama Y., Shinomura Y., Role of matrix metalloproteinase-7 (matrilysin) in human cancer invasion, apoptosis, growth, and angiogenesis. *Exp. Biol. Med.*, 2006, 231, 20–27.
  22. Jaszewska E., Kośmider A., Kiss A.K., Naruszewicz M., Pro-oxidative and pro-apoptotic action of defatted seeds of *Oenothera paradoxa* on human skin melanoma cells. *J. Agric. Food Chem.*, 2009, 57, 8282–8289.
  23. Jaszewska E., Kosmider A., Kiss A.K., Naruszewicz M., *Oenothera paradoxa* defatted seeds extract containing pentagalloylglucose and procyanidins potentiates the cytotoxicity of vincristine. *J. Physiol. Pharmacol.*, 2010, 61, 637–643.
  24. Jemal A., Bray F., Center M.M., Ferlay J., Ward E., Forman D., Global cancer statistics. *CA Cancer J. Clin.*, 2011, 61, 69–90.
  25. Kessenbrock K., Plaks V., Werb Z., Matrix metalloproteinases: regulators of the tumor microenvironment. *Cell*, 2010, 141, 52–67.
  26. Kim M., Murakami A., Ohigashi H., Modifying effects of dietary factors on (-)-epigallocatechin-3-gallate-induced pro-matrix metalloproteinase-7 production in HT-29 human colorectal cancer cells. *Biosci. Biotechnol. Biochem.*, 2007, 71, 2442–2450.
  27. Kiss A.K., Derwińska M., Dawidowska A., Naruszewicz M., Novel biological properties of *Oenothera paradoxa* defatted seed extracts: effects on metalloproteinase activity. *J. Agric. Food Chem.*, 2008, 56, 7845–7852.
  28. Kiss A.K., Kapłon-Cieślicka A., Filipiak K.J., Opolski G., Naruszewicz M., *Ex vivo* effects of an *Oenothera paradoxa* extract on the reactive oxygen species generation and neutral endopeptidase activity in neutrophils from patients after acute myocardial infarction. *Phytother. Res.*, 2012, 26, 482–487.
  29. Kołomecki K., Hamowanie aktywności metaloproteinaz – możliwości zastosowania klinicznego. *Onkol. Pol.*, 2000, 3, 163–167 (in Polish).
  30. Kumar A., Naidu P.S., Seghal N., Padi S.S.V., Neuroprotective effects of resveratrol against intracerebroventricular colchicine-induced cognitive impairment and oxidative stress in rats. *Pharmacology*, 2007, 79, 17–26.
  31. Kupai K., Szucs G., Cseh S., Hajdu I., Csonka C., Csont T., Ferdinandy P., Matrix metalloproteinase activity assays: Importance of zymography. *J. Pharmacol. Toxicol. Methods*, 2010, 61, 205–209.
  32. Labrie M., St-Pierre Y., Epigenetic regulation of mmp-9 gene expression. *Cell Mol. Life Sci.*, 2013, 70, 3109–3124.
  33. Lamy S., Blanchette M., Michaud-Levesque J., Laffleur R., Durocher Y., Moghrabi A., Barrette S., Gingras D., Béliveau R., Delphinidin, a dietary anthocyanidin, inhibits vascular endothelial growth factor receptor-2 phosphorylation. *Carcinogenesis*, 2006, 27, 989–996.
  34. Letchoumy P.V., Mohan K.V., Prathiba D., Hara Y., Nagini S., Comparative evaluation of antiproliferative, antiangiogenic and apoptosis inducing potential of black tea polyphenols in the hamster buccal pouch carcinogenesis model. *J. Carcinog.*, 2007, 6, 19.
  35. Lewandowska U., Szewczyk K., Owczarek K., Hrabec Z., Podśędek A., Sosnowska D., Hrabec E., Procyanidins from evening primrose (*Oenothera paradoxa*) defatted seeds inhibit invasiveness of breast cancer cells and modulate the expression of selected genes involved in angiogenesis, metastasis, and apoptosis. *Nutr. Cancer*, 2013, 65, 1219–1231.
  36. Li X.L., Sun S., Du G.J., Qi L.W., Williams S., Wang C.Z., Yuan C.S., Effects of *Oplopanax horridus* on human colorectal cancer cells. *Anticancer Res.*, 2010, 30, 295–302.
  37. Lubbe W.J., Zhou Z.Y., Fu W., Zuzga D., Schulz S., Fridman R., Muschel R.J., Waldman S.A., Pitari G.M., Tumor epithelial cell matrix metalloproteinase 9 is a target for antimetastatic therapy in colorectal cancer. *Clin. Cancer Res.*, 2006, 12, 1876–1882.
  38. Mertens-Talcott S.U., Percival S.S., Ellagic acid and quercetin interact synergistically with resveratrol in the induction of apoptosis and cause transient cell cycle arrest in human leukemia cells. *Cancer Lett.*, 2005, 218, 141–151.
  39. Naczki M., Amarowicz R., Zadernowski R., Shahidi F., Protein precipitating capacity of condensed tannins of beach pea, canola hulls, evening primrose and faba bean. *Food Chem.*, 2001, 73, 467–471.
  40. Nakamura T., Kuwai T., Kim J.S., Fan D., Kim S.J., Fidler I.J., Stromal metalloproteinase-9 is essential to angiogenesis and progressive growth of orthotopic human pancreatic cancer in parabiont nude mice. *Neoplasia*, 2007, 9, 979–986.
  41. Noël A., Jost M., Maquoi E., [Matrix metalloproteinases at cancer tumor-host interface. Semin.](#) *Cell Dev. Biol.*, 2008, 19, 52–60.
  42. Ogasawara M., Matsunaga T., Suzuki H., Differential effects of antioxidants on the *in vitro* invasion, growth and lung metastasis of murine colon cancer cells. *Biol. Pharm. Bull.*, 2007, 30, 200–204.
  43. Peiretti P.G., Palmegiano G.B., Masoero G., Chemical composition, organic matter digestibility and fatty acid content of evening primrose (*Oenothera paradoxa*) during its growth cycle. *Anim. Feed Sci. Tech.*, 2004, 116, 293–299.
  44. Peri C., Pompei C., An assay of different phenolic fractions in wines. *Amer. J. Enol. Vitic.*, 1971, 22, 55–58.
  45. Poincloux R., Lizárraga F., Chavier P., Matrix invasion by tumour cells: a focus on MT1-MMP trafficking to invadopodia. *J. Cell Sci.*, 2009, 122, 3015–3024.
  46. Ra H.J., Parks W.C., Control of matrix metalloproteinase catalytic activity. *Matrix Biol.*, 2007, 26, 587–596.
  47. Robaszkiewicz A., Balcerczyk A., Bartosz G., Antioxidative and prooxidative effects of quercetin on A549 cells. *Cell Biol. Int.*, 2007, 31, 1245–1250.
  48. Rösch D., Bergmann M., Knorr D., Kroh L.W., Structure – antioxidant efficiency relationships of phenolic compounds and their contribution to the antioxidant activity of sea buckthorn juice. *J. Agric. Food Chem.*, 2003, 51, 4233–4239.
  49. Stręć M., Goriach S., Podśędek A., Sosnowska D., Koziolkiewicz M., Hrabec Z., Hrabec E., Procyanidin oligomers from Japanese quince (*Chaenomeles japonica*) fruit inhibit activity of MMP-2 and MMP-9 metalloproteinases. *J. Agric. Food Chem.*, 2007, 55, 6447–6452.
  50. Swain T., Hillis W.E., The phenolic constituents of *Prunus domestica* I. *J. Sci. Food Agric.*, 1959, 10, 63–68.

51. Toegel S., Wu S.Q., Otero M., Goldring M.B., Leelapornpisid P., Chiari C., Kolb A., Unger F.M., Windhager R., Viernstein H., *Caesalpinia sappan* extract inhibits IL1 $\beta$ -mediated overexpression of matrix metalloproteinases in human chondrocytes. *Genes Nutr.*, 2012, 7, 307–318.
52. Yang J., Wei D., Liu J., Repressions of MMP-9 expression and NF-kappa B localization are involved in inhibition of lung carcinoma 95-D cell invasion by (-)-epigallocatechin-3-gallate. *Biomed. Pharmacother.*, 2005, 59, 98–103.
53. Yun J.H., Pang E.K., Kim C.S., Yoo Y.J., Cho K.S., Chai J.K., Kim C.K., Choi S.H., Inhibitory effects of green tea polyphenol (-)-epigallocatechin gallate on the expression of matrix metalloproteinase-9 and on the formation of osteoclasts. *J. Periodontal Res.*, 2004, 39, 300–307.

Submitted: 21 August 2013. Revised: 30 October 2013 and 20 January 2014. Accepted: 3 February 2014. Published on-line: 10 June 2014.